

兔抗 CCT6A 多克隆抗体

中文名称：兔抗 CCT6A 多克隆抗体

英文名称： Anti-CCT6A rabbit polyclonal antibody

别名： CCT6; Cctz; HTR3; TCPZ; TCP20; MoDP-2; TTCP20; CCT-zeta; CCT-zeta-1; TCP-1-zeta

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： CCT6A

反应种属： Human

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

Background:

The protein encoded by this gene is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP 1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Alternate transcriptional splice variants of this gene, encoding different isoforms, have been characterized. In addition, several pseudogenes of this gene have been located

| | |
|---------------------------------|---|
| | d. |
| Applications: | WB, IHC |
| Name of antibody: | CCT6A |
| Immunogen: | Synthesized peptide derived from internal of human CCT6A. |
| Full name: | chaperonin containing TCP1, subunit 6A (zeta 1) |
| Synonyms : | CCT6; Cctz; HTR3; TCPZ; TCP20; MoDP-2; TTCP20; CCT-zeta; CCT-zeta-1; TCP-1-zeta |
| SwissProt: | P40227 |
| IHC positive control: | Human breast carcinoma tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 58 kDa |
| WB Positive control: | HepG2 cells lysate |
| WB Recommended dilution: | 500-3000 |



