

兔抗 TAP2 多克隆抗体

- 中文名称：兔抗 TAP2 多克隆抗体
- 英文名称：Anti-TAP2 rabbit polyclonal antibody
- 别名：APT2; PSF2; ABC18; ABCB3; PSF-2; RING11; D6S217E
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：TAP2
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

| | |
|----------------------|--|
| Background: | The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. This gene is located 7 kb telomeric to gene family member ABCB2. |
| Applications: | ELISA, WB |

| | |
|------------------------------------|---|
| Name of antibody: | TAP2 |
| Immunogen: | Synthetic peptide of human TAP2 |
| Full name: | transporter 2, ATP-binding cassette, sub-family B (MDR/TAP) |
| Synonyms : | APT2; PSF2; ABC18; ABCB3; PSF-2; RING11; D6S217E |
| SwissProt: | Q03519 |
| ELISA Recommended dilution: | 1000-2000 |
| WB Predicted band size: | 76 kDa |
| WB Positive control: | Jurkat cells |
| WB Recommended dilution: | 200-1000 |

